Pancreas disease – factors influencing transmission and infection

Sonal Patel Institute of Marine Research Vaxxinova Norway

PD trination 2018




Research council of Norway: Toppforsk research Grant 224885/E40, MitSAV

Ministry of trade, industry and fisheries through Aquaculture program, IMR

NORAD

Projects: MitSAV and PATHOS

- 1. Understand the effects of physiological and environmental factors on SAV infectivity, latency, and survival, and how these relate to the development of the clinical symptoms of PD
- 2. Develop methods and tools to better understand the immune response of Atlantic salmon to SAV3




Objectives

- 1. <u>Develop tools and assays esp SAV challenge model in post-smolt</u> stages
- 2. Characterise the effect of <u>sub-optimal physiological and</u> <u>environmental parameters</u> on host, pathogen and host-pathogen interaction
- 3. Characterise the role of <u>specific immune cells</u>, and immune <u>response</u> during a SAV infection
- 4. Investigate the SAV susceptibility and immune response in both <u>diploid and triploid Atlantic salmon</u>
- 5. Study correlation between SAV3 infection and microbiome of salmon

Factors influencing salmon-SAV interaction <u>Salmon:</u>

- Physiology
- Innate and adaptive immune responses
- Genetics and Ploidy



Environment/other factors:

- Time
- Temperature
- Organic load
- Oxygen
- UV radiation
- Microbiome

<u>SAV:</u>

- Sub-type and isolate
- Virulence
- Infectivity and dose
- Host specificity
- Survival in and outside host


Objective

- 1. <u>Develop tools and assays</u> esp SAV challenge model in post-smolt stages
 - Establish bath challenge model with SAV3 in post-smolts
 - Optimise SAV analysis method for seawater

Bath challenge model

qPCR and TCID50

Natural infection route Defined time of infection • Dose quantitated by RT-




Shedder fish got sick and shed virus into

Remove shedders from seawater containing virus for bath challenge

Add naïve fish into seawater containing

Optimised method for quntitating SAV in water




Objective

2. Characterise the effect of <u>sub-optimal physiological and</u> <u>environmental parameters</u> on host, pathogen and host-pathogen interaction

- Susceptibility of salmon at two different time-points post sea water transfer
- Impact of viral dose on infection dynamics
- Survival of SAV at different temperatures and in different "medium"

Susceptibility to SAV3




Jarungsriapisit et al, 2016a

Viral shedding in tank-water – RT-qPCR


Shedding from shedders

Shedding from fish bath challenged in shedder water



Jarungsriapisit et al, 2016a


Time for adaptation to seawater - susceptibility



Phase B: Lower prevalance, Lower viral loads and lower shedding


Jarungsriapisit et al, 2016a

Dose study - Experimental setup



Jarungsriapisit et al, 2016b


Shedding in water



Days post infection


Jarungsriapisit et al, 2016b

Prevalance




Dose study

Viral loads



Jarungsriapisit et al, 2016b

Survival of SAV3




Jarungsriapisit et al, in prep

Objective

3. Characterise the role of <u>specific immune cells</u>, and immune <u>response</u> during a SAV infection


- Gene regulation of innate and adaptive genes in the salmon at two phases
- Immune cells in SAV affected tissues

Innate immune genes in head kidney of non-infected fish




Moore et al, 2017 and 2018

Innate immune genes in head kidney of SAV3 infected salmon




Moore *et al.*, 2017 and 2018

Affected tissues -histology



Jarungsriapisit et al, submitted




Cells in affected tissues -IHC

Scores

0

2


3



Objective

4. Investigate the SAV susceptibility and immune response in both <u>diploid and triploid</u> Atlantic salmon

Atlantic salmon




Presentation by Noelia Nunez Ortiz

Sub-topic

5. Study correlation between SAV3 infection and microbiome of salmon

- Co-relation between microbiome and dose of SAV3 in salmon skin
- Compare gill microbiome in triploid and diploid salmon


Microbiome from Dose study- Skin



SAV3 infection results in losses of beneficial bacteria in Atlantic salmon skin

Reid et al 2017


Skin dysbiosis - micorbiome



Factors influencing salmon-SAV interaction - summary

Salmon:

- Physiology
- Innate and adaptive immune responses
- Genetics and Ploidy



Environment/other factors:

- Time
- Temperature
- Organic load
- Oxygen
- UV radiation
- Microbiome

<u>SAV:</u>

- Sub-type and isolate
- Virulence
- Infectivity and dose
- Host specificity
- Survival in and outside host

Thanks to..

Institute of Marine Research:

Postdoc Lindsey Moore, Noelia Nunez Ortiz PhD: Jiraporn Jarungsriapisit Stig Mæhle, Ann Cathrine Bårdsgjære Einen, Joachim Nordbø, Ingrid U. Fiksdal, Cecilie Isaachsen, Miriam Castillo Furne, Cecilie Skår, Ivar Helge Matre Dr. Geir Lasse Taranger, Dr. Per Gunnar Fjelldal, Dr. H. Craig Morton

University of Bergen: Professor Sigurd Stefansson Uni Research Bergen: Dr. Tom Ole Nilsen Norwegian School of Veterinary Science: Professor Øystein Evensen Friedrich-Loeffler-Institut, Germany: Dr. Bernd Köllner University of Aberdeen, UK: Dr. Samuel Martin, Professor Chris Secombes University of New Mexico, USA: Dr. Irene Salinas and her lab Vaxxinova Norway: Arild Tangerås

Petter Frost, MSD animal health Norway